

- are improving: Results from a population-based registry. *J Vasc Surg* 2006;44:79-85.
21. LaMuraglia GM, Brewster DC, Moncure AC, Dorer DJ, Stoner MC, Trehan SK, et al. Carotid endarterectomy at the millennium: what interventional therapy must match. *Ann Surg* 2004;240:535-46.
 22. Khuri SF, Daley J, Henderson W, Barbour GJ, Lowry P, Irvin G, et al. National Veterans Administration Surgical Risk Study: risk adjustment for the comparative assessment of the quality of surgical care. *J Am Coll Surg* 1995;180:519-31.
 23. Fink AS, Campbell DA Jr, Mentzer RM Jr, Henderson WG, Daley J, Bannister J, et al. The National Surgical Quality Improvement Program in non-veterans administration hospitals: initial demonstration of feasibility. *Ann Surg* 2002;236:344-53.
 24. CAVATAS investigators. Endovascular versus surgical treatment in patients with carotid stenosis in the carotid and vertebral artery transluminal angioplasty study (CAVATAS): a randomised trial. *Lancet* 2001;357:1729-37.
 25. Alberts MJ. Results of a multicenter prospective randomized trial of carotid artery stenting vs carotid endarterectomy [abstract]. *Stroke* 2001;32:325-d.
 26. SPACE collaborative group, Ringleb PA, Allenberg J, Bruckmann H, Eckstein HH, Fraedrich G, Hartmann M, et al. 30 day results from the SPACE trial of stent-protected angioplasty versus carotid endarterectomy in symptomatic patients: a randomised non-inferiority trial. *Lancet* 2006;368:1239-47.
 27. Mas JL, Chatellier G, Beyssen B, Branchereau A, Moulin T, Becquemin JP, et al; EVA-3S investigators. Endarterectomy versus stenting in patients with symptomatic severe carotid stenosis. *N Engl J Med* 2006;355:1660-71.
 28. Hobson RW II, Howard VJ, Roubin GS, Brott TG, Ferguson RD, Popma JJ, et al, for the CREST Investigators. Carotid artery stenting is associated with increased complications in octogenarians: 30-day stroke and death rates in the CREST lead-in phase. *J Vasc Surg* 2004;40:1106-11.
 29. Gray WA, Hopkins LN, Yadav S, Davis T, Wholey M, Atkinson R, et al; ARChER trial collaborators. Protected carotid stenting in high-surgical-risk patients: the ARChER results. *J Vasc Surg* 2006;44:258-69.
 30. Iyer SS, White CJ, Hopkins LN, Katzen BT, Safian R, Wholey MH, et al; BEACH investigators. Carotid artery revascularization in high-surgical-risk patients using the Carotid WALLSTENT and FilterWire EX/EZ: 1-year outcomes in the BEACH Pivotal Group. *J Am Coll Cardiol* 2008;51:427-34.
 31. Flanigan DP, Flanigan ME, Dorne AL, Harward TRS, Razavi MK, Ballard JL. Long-term results of 442 consecutive, standardized carotid endarterectomy procedures in standard-risk and high-risk patients. *J Vasc Surg* 2007;46:876-82.
 32. Mozes G, Sullivan TM, Torres-Russotto DR, Bower TC, Hoskin TL, Sampaio SM, et al. Carotid endarterectomy in SAPHIRE-eligible high-risk patients: Implications for selecting patients for carotid angioplasty and stenting. *J Vasc Surg* 2004;39:958-66.
 33. Boules TN, Proctor MC, Aref A, Upchurch GR Jr, Stanley JC, Henke PK. Carotid endarterectomy remains the standard of care, even in high-risk surgical patients. *Ann Surg* 2005;241:356-63.
 34. Illig KA, Zhang R, Tanski W, Benesch C, Sternbach Y, Green RM. Is the rationale for carotid angioplasty and stenting in patients excluded from NASCET/ACAS or eligible for ARChER justified? *J Vasc Surg* 2003;37:575-81.
 35. Lepore MR, Sternbergh WC III, Salartash K, Tonnessen B, Money SR. Influence of NASCET/ACAS trial eligibility on outcome after carotid endarterectomy. *J Vasc Surg* 2001;34:581-6.
 36. Miller MT, Comerota AJ, Tzilinis A, Daoud Y, Hammerling J. Carotid endarterectomy in octogenarians: Does increased age indicate "high risk?" *J Vasc Surg* 2005;41:231-7.
 37. Sayeed S, Stanziale SF, Wholey MH, Makaroun MS. Angiographic lesion characteristics can predict adverse outcomes after carotid artery stenting. *J Vasc Surg* 2008;47:81-7.
 38. Lam RC, Lin SC, DeRubertis B, Hyncek R, Kent KC, Faries PL. The impact of increasing age on anatomic factors affectin carotid angioplasty and stenting. *J Vasc Surg* 2007;45:875-80.
 39. Stanziale SF, Marone LK, Boules TN, Brimmer JA, Hill K, Makaroun MS, et al. Carotid artery stenting in octogenarians is associated with increased adverse outcomes. *J Vasc Surg* 2006;43:297-304.
 40. Krejza J, Arkuszewski M, Kasner SE, Weigle J, Ustymowicz A, Hurst RW, et al. Carotid artery diameter in men and women and the relation to body and neck size. *Stroke* 2006;37:1103-5.
 41. Messe SR, Kasner SE, Mehta Z, Warlow CP, Rothwell PM, European Carotid Surgery Trialists. Effect of body size on operative risk of carotid endarterectomy. *J Neurol Neurosurg Psychiatry* 2004;75:1759-61.
 42. Rothwell PM, Eliasziw M, Gutnikov SA, Warlow CP, Barnett HJ; Carotid Endarterectomy Trialists Collaboration. Endarterectomy for symptomatic carotid stenosis in relation to clinical subgroups and timing of surgery. *Lancet* 2004;363:915-24.
 43. Alamowitch S, Eliasziw M, Barnett HJ. The risk and benefit of endarterectomy in women with symptomatic internal carotid artery disease. *Stroke* 2005;36:27-31.
 44. Rothwell PM, Goldstein LB. Carotid endarterectomy for asymptomatic carotid stenosis: asymptomatic carotid surgery trial. *Stroke* 2004;35:2425-7.

Submitted Jun 29, 2008; accepted Sep 11, 2008.

DISCUSSION

Dr Amy Reed (Cincinnati, OH). Were you able to look at the asymptomatic vs symptomatic indication for carotid endarterectomy and particularly look at how those high-risk patients did? There have been studies that have shown that that is a particular area where patients don't do well and whether or not best medical management might be better.

Dr Jeanwan Kang. The information that is available in the NSQIP database is a history of previous neurologic event. What we don't know is the laterality or the timing of that event, so we are not able to determine who underwent the surgery for symptomatic vs asymptomatic disease.

What we are able to determine from the database is a group of patient population who, for sure, is asymptomatic, mainly those with absolutely no previous neurologic event, and in those patients there was a slightly lower rate of stroke death rate, however it was not statistically significant.

Dr Hasan Dosluoglu (Buffalo, NY). I enjoyed the talk and a wealth of information from the private NSQIP data is going to start coming out. I am surprised that 80% of the patients had actually general anesthesia. As a local anesthesia user for this

procedure, I am surprised at that. Did the high-risk patients have different outcomes in those who had a general anesthesia vs local anesthesia? And second, maybe the high-risk patients were not too high risk, because they could undergo general anesthesia in those who had general anesthesia.

Dr Kang. To answer your first question, because the percentage of patients undergoing local anesthesia was less than 20%, that came out to about a little over 600 patients undergoing carotid endarterectomy under local anesthesia, so any sort of subanalysis became meaningless because the numbers were so low.

The criteria for high risk were the same as that defined by the SAPHIRE study and there were about 30% of patients who met these criteria. In addition, ASA [American Society of Anesthesiologists] class for each patient is available in the NSQIP database. I have not shown these data here, but patients who were ASA class 3 or 4 made up majority of the patient population.

Dr Anton Sidawy (Washington, DC). We analyzed the VA NSQIP data. There were over 20,000 carotid endarterectomies. Although there was no significant difference in neurologic outcomes based on different levels of renal insufficiency, 30-day

mortality and pulmonary, cardiac, and infectious complications were significantly increased. Have you adjusted your data for renal failure?

Dr Kang. It certainly was. For stroke, renal failure was not one of the variables that came out to be significant on univariate analysis. And we looked at renal failure in many different ways. We looked at it as a categorical variable as well as continuous variable. We also calculated GFR and none of that made a difference in

terms of univariate analysis for stroke. It was, however, one of the variables that came out to be significant for death. However, on multivariate analysis, it was not one of the independent predictors of death.

Having said that, the number of deaths that were studied in our study were only 26; and so therefore, we could only say that two of the highest, most significant variables on this multivariate analysis were significant predictors of death.